

- |                               |                                |                                    |                                     |
|-------------------------------|--------------------------------|------------------------------------|-------------------------------------|
| <input type="checkbox"/> HLAS | <input type="checkbox"/> HSA   | <input type="checkbox"/> CHSLD DBV | <input type="checkbox"/> CHSLD GD   |
| <input type="checkbox"/> HGL  | <input type="checkbox"/> IUSMD | <input type="checkbox"/> CHSLD DOR | <input type="checkbox"/> CHSLD NP   |
| <input type="checkbox"/> CHSM | <input type="checkbox"/> CLSC  | <input type="checkbox"/> CHSLD LAS | <input type="checkbox"/> CHSLD LACH |

N° dossier / Chart n°	DDN / DOB	Sexe / Sex
Nom / Name	Prénom / First Name	
Nom de la mère / Name of mother		
Adresse / Address		
Tél. / Tel.		
N° assurance maladie / Medicare Card N°		Expiration

Allergie(s): \_\_\_\_\_ Height: \_\_\_\_\_ Weight: \_\_\_\_\_

Prescriber's initials	<b>Enhanced recovery after surgery (ERAS) - Colorectal Per-operative prescriptions</b>
	<b><u>Anti-emetic prophylaxis</u></b>
	Dexamethasone <input type="checkbox"/> 4 mg <b>OR</b> <input type="checkbox"/> 6 mg IV X 1 dose
	Ondansetron 4 mg IV X 1 dose
	Other: _____
	<b><u>Antimicrobial prophylaxis</u></b>
	<b>Administer 0-60 minutes before incision:</b>
	Cefazoline 2 g IV X 1 dose over 5 min + Metronidazole 500 mg IV X 1 dose over 20 min <b>OR</b>
	Cefazoline 3 g IV X 1 dose over 5 min + Metronidazole 500 mg IV X 1 dose over 20 min ( <b>if patient ≥ 120 kg</b> )
	<i>If anaphylactic allergy to penicillin (infectious diseases consultation recommended):</i>
	Metronidazole 500 mg IV X 1 dose over 20 min <b>AND</b>
	Ciprofloxacin 400 mg IV X 1 dose over 30 min
	<b><u>Administer a second dose of antimicrobial prophylaxis (in patients with normal renal function) if the patient has lost more than 1500 mL of blood AND/OR</u></b>
	Cefazoline 2 g IV X 1 dose over 5 min (if surgery lasts more than 4 hours)
	Metronidazole 500 mg IV X 1 dose over 20 min (if surgery lasts more than 6 hours)
	<b><u>Anaesthetic approaches reducing the use of opioids</u></b>
	✓ Use an anaesthetic approach that reduces the use of opioids and clearly document this in the anaesthesia sheet.
	<b><u>Maintenance of normothermia</u></b>
	✓ Aim to maintain body temperature between ≥ 36.0°C and < 38.0°C throughout the procedure and document.

GRM 4200043369

Prescriber: Name (block letters)	Signature	License No	Date (YYYY/MM/DD)	Hour